arXiv:math/0512358v2 [math.CA] 16 Jan 2006

Discrete Painlevé equations for recurrence coefficients of orthogonal polynomials

Walter Van Assche *
Katholieke Universiteit Leuven, Belgium

1 Introduction

Orthonormal polynomials on the real line are defined by the orthogonality conditions

$$\int_{\mathbb{R}} p_n(x)p_m(x)\,d\mu(x) = \delta_{m,n}, \qquad m,n \geq 0, \tag{1.1}$$

where μ is a positive measure on the real line for which all the moments exist and $p_n(x) = \gamma_n x^n + \cdots$, with positive leading coefficient $\gamma_n > 0$. A family of orthonormal polynomials always satisfies a three-term recurrence relation of the form

$$xp_n(x) = a_{n+1}p_{n+1}(x) + b_n p_n(x) + a_n p_{n-1}(x), \qquad n \geq 0, \tag{1.2}$$

with $p_{-1} = 0$ and

$$p_0 = \gamma_0 = \left(\int d\mu(x)\right)^{-1/2}. \tag{1.3}$$

Comparing the leading coefficients in (1.2) gives

$$a_{n+1} = \frac{\gamma_n}{\gamma_{n+1}} > 0, \tag{1.4}$$

and computing the Fourier coefficients of $xp_n(x)$ in (1.2) gives

$$a_n = \int_{\mathbb{R}} xp_n(x)p_{n-1}(x)\,d\mu(x), \tag{1.5}$$

$$b_n = \int_{\mathbb{R}} xp_n^2(x)\,d\mu(x). \tag{1.6}$$

The converse statement is also true and is known as *the spectral theorem for orthogonal polynomials*: if a family of polynomials satisfies a three-term recurrence relation of the form (1.2), with $a_n > 0$ and $b_n \in \mathbb{R}$ and initial conditions $p_0 = 1$ and $p_{-1} = 0$, then there exists a *probability measure* μ on the real line such that these polynomials are orthonormal polynomials satisfying (1.1). This gives rise to two important problems:

---

*Research supported by INTAS Research Network NeCCA (03-51-6637), FWO projects G.0184.02 and G.0455.04, and OT/04/21 of the Research Counsel of K.U.Leuven.

Problem 1. Suppose the measure μ is known. What can be said about the recurrence coefficients $\{a_n : n = 1, 2, 3, \ldots\}$ and $\{b_n : n = 0, 1, 2, \ldots\}$? This is known as the *direct problem* for orthogonal polynomials.

Problem 2. Suppose the recurrence coefficients $\{a_{n+1}, b_n : n = 0, 1, 2, \ldots\}$ are known. What can be said about the orthogonality measure μ? This is known as the *inverse problem* for orthogonal polynomials.

The recurrence coefficients are usually collected in a tridiagonal matrix of the form

$$J = \begin{pmatrix} b_0 & a_1 & & & \\ a_1 & b_1 & a_2 & & \\ & a_2 & b_2 & a_3 & \\ & & a_3 & b_3 & \ddots \\ & & & \ddots & \ddots \end{pmatrix} \tag{1.7}$$

which acts as an operator on (a subset of) $\ell_2(\mathbb{N})$ and which is known as the Jacobi matrix or Jacobi operator. If J is self-adjoint, then the spectral measure for J is precisely the orthogonality measure μ. Hence problem 1 corresponds to the inverse problem for the Jacobi matrix J and problem 2 corresponds to the direct problem for J.

In the present paper we will study problem 1 for a few special cases. In Section 2 we study measures on the real line with an exponential weight function of the form $d\mu(x) = |x|^\rho \exp(-|x|^m)\, dx$, which are known as Freud weights, named after Géza Freud who studied them in the 1970's. It will be shown that the recurrence coefficients a_n satisfy a non-linear recurrence relation which corresponds to the discrete Painlevé I equation and its hierarchy. In Section 3 we will study a family of orthogonal polynomials on the unit circle. We will first give some background on orthogonal polynomials on the unit circle and the corresponding recurrence relations. We will study the weight function $w(\theta) = \exp(\lambda \cos\theta)$, and it will be shown that the recurrence coefficients satisfy a non-linear recurrence relation which corresponds to discrete Painlevé II. These orthogonal polynomials play an important role in random unitary matrices and combinatorial problems for random permutations. In Section 4 we will study certain discrete orthogonal polynomials related to Charlier polynomials. The recurrence coefficients a_n and b_n are shown to satisfy a system of non-linear recurrence relations which are again related to the discrete Painlevé II equation. Finally, in Section 5 we consider certain q-orthogonal polynomials which are q-analogs of the Freud weight. We will show that the recurrence coefficients satisfy a q-deformed Painlevé I equation. Most of the material in this paper is not new: the recurrence relations in Section 2 were already obtained by Freud in [7] and are known as *Freud equations* in the field of orthogonal polynomials. It was Magnus [17] who made the connection with the discrete Painlevé equation I. The recurrence relation in Section 3 was found by Periwal and Shevitz [20] (see also Hisakado [9], Tracy and Widom [23]; Baik [1] used the Riemann-Hilbert approach to obtain the Painlevé equation). The recurrence relations in Section 4 were obtained by Van Assche and Foupouagnigni in [25]. The results in Section 5 were first obtained by Nijhoff [19]. We hope that bringing together these various orthogonal polynomials and the corresponding discrete Painlevé equations will be illuminating and encourage researchers in the field of orthogonal polynomials and researchers in integrable difference equations to talk to each other, and that the interaction between both areas of mathematics will shed some extra light on either subject.

2 Freud weights

Freud weights are exponential weights on the real line $(-\infty,\infty)$ of the form

$$w_\rho(x) = |x|^\rho \exp(-|x|^m), \qquad \rho > -1, m > 0.$$

They were considered by Géza Freud in his 1976 paper [7], where he gave a recurrence relation for the recurrence coefficients a_n when $m = 2, 4, 6$. For these cases Freud found the asymptotic behavior of the recurrence coefficients a_n and he formulated a conjecture for this asymptotic behavior for every $m > 0$. This conjecture really started the analysis of general orthogonal polynomials on unbounded sets of the real line, since before Freud's work only very special cases such as the Hermite polynomials and the Laguerre polynomials were studied in detail. In this section I will repeat Freud's analysis of the recurrence coefficients of Freud weights, make the connection with discrete Painlevé equations (which was not known to Freud but first pointed out by Magnus in [17]), and point out what has been done after Freud. Observe that Freud weights are symmetric, i.e., $w_\rho(-x) = w_\rho(x)$, which implies that $b_n = 0$ for $n \geq 0$.

2.1 Generalized Hermite polynomials

The case $m = 2$ corresponds to generalized Hermite polynomials (and $\rho = 0$ are the Hermite polynomials). Generalized Hermite polynomials were already investigated by Chihara in [3] (see also [4]). The weight $w_0(x) = \exp(-x^2)$ satisfies the first order differential equation

$$[w_0(x)]' = -2xw_0(x) \tag{2.1}$$

which is the *Pearson equation* for the Hermite weight. In general a weight w satisfying a Pearson equation of the form $[\sigma(x)w(x)]' = \tau(x)w(x)$, with σ a polynomial of degree at most two and τ a polynomial of degree one, is called a classical weight. The weight functions for Hermite polynomials, Laguerre polynomials, and Jacobi polynomials are the classical weights for σ of degree zero, one and two, respectively. Bessel polynomials appear for $\sigma(x) = x^2$, but they are not orthogonal on the real line with respect to a positive measure. Freud's idea was to compute the integral

$$\int_{-\infty}^{\infty} (p_n(x)p_{n-1}(x)|x|^\rho)'\, w_0(x)\, dx \tag{2.2}$$

in two different ways. The first way is simply working out the derivative in the integrand and to use the orthogonality to evaluate the resulting terms. This gives

$$\begin{aligned}
\int_{-\infty}^{\infty} (p_n(x)p_{n-1}(x)|x|^\rho)'\, w_0(x)\, dx &= \int_{-\infty}^{\infty} p_n'(x)p_{n-1}(x)|x|^\rho w_0(x)\, dx && (2.3)\\
&+ \int_{-\infty}^{\infty} p_n(x)p_{n-1}'(x)|x|^\rho w_0(x)\, dx && (2.4)\\
&+ \rho \int_{-\infty}^{\infty} \frac{p_n(x)p_{n-1}(x)}{x}|x|^\rho w_0(x)\, dx. && (2.5)
\end{aligned}$$

For the right hand side in (2.3) we use the fact that

$$\begin{aligned}
p_n'(x) &= n\gamma_n x^{n-1} + \text{ lower order terms}\\
&= n\frac{\gamma_n}{\gamma_{n-1}}p_{n-1}(x) + \text{ lower degree terms.}
\end{aligned}$$

This gives

$$\int_{-\infty}^{\infty} p_n'(x)p_{n-1}(x)|x|^\rho w_0(x)\, dx = n\frac{\gamma_n}{\gamma_{n-1}}.$$

For the integral in (2.4) we see that $p_{n-1}'(x)$ is a polynomial of degree $n-2$ and hence by orthogonality the integral vanishes. For the integral in (2.5) we use the fact that the weight function w_ρ is even, i.e., $w_\rho(-x) = w_\rho(x)$, which implies that $p_n(-x) = (-1)^n p_n(x)$. This means that $p_n(x)/x$ is a polynomial of degree $n-1$ when n is odd and $p_{n-1}(x)/x$ is a polynomial of degree $n-2$ when n is even. Hence when n is even the integral in (2.5) vanishes, and when n is odd we have

$$\frac{p_n(x)}{x} = \frac{\gamma_n}{\gamma_{n-1}} p_{n-1}(x) + \text{ lower degree terms,}$$

so that

$$\rho \int_{-\infty}^{\infty} \frac{p_n(x)p_{n-1}(x)}{x}|x|^\rho w_0(x)\, dx = \rho\frac{\gamma_n}{\gamma_{n-1}}\Delta_n$$

where

$$\Delta_n = \begin{cases} 0 & \text{if } n \text{ is even,} \\ 1 & \text{if } n \text{ is odd.} \end{cases}$$

Combining these results and using the expression $a_n = \gamma_{n-1}/\gamma_n$ gives

$$\int_{-\infty}^{\infty} \left(p_n(x)p_{n-1}(x)|x|^\rho\right)' w_0(x)\, dx = \frac{n+\rho\Delta_n}{a_n}. \tag{2.6}$$

Observe that this holds whenever w is a symmetric weight on the real line. A second way to compute the integral in (2.2) is to use integration by parts, combined with Pearson's equation (2.1) for the weight. This gives

$$\begin{aligned} \int_{-\infty}^{\infty} \left(p_n(x)p_{n-1}(x)|x|^\rho\right)' w_0(x)\, dx &= -\int_{-\infty}^{\infty} p_n(x)p_{n-1}(x)|x|^\rho w_0'(x)\, dx \\ &= 2\int_{-\infty}^{\infty} xp_n(x)p_{n-1}(x)|x|^\rho w_0(x)\, dx \\ &= 2a_n. \end{aligned} \tag{2.7}$$

Combining (2.6) and (2.7) then gives

$$a_n^2 = \frac{n+\rho\Delta_n}{2} \tag{2.8}$$

so that $a_n = \sqrt{n+\rho\Delta_n}/\sqrt{2}$. For $\rho = 0$, which are the Hermite polynomials, one has $a_n = \sqrt{n/2}$. For generalized Hermite polynomials one has

$$\lim_{n\to\infty} \frac{a_n}{\sqrt{n}} = \frac{1}{\sqrt{2}}. \tag{2.9}$$

2.2 Freud weight $m = 4$

Let us now consider the weight $w_\rho(x) = |x|^\rho \exp(-x^4)$, where $\rho > -1$. The Pearson equation for the weight $w_0(x) = \exp(-x^4)$ is

$$[w_0(x)]' = -4x^3 w_0(x),$$

which is a first order linear differential equation with polynomial coefficients, but since the polynomial coefficient is now of degree 3, the weight w_0 is no longer a classical weight but a semi-classical one. The equation (2.6) remains valid for this Freud weight, but integration by parts give a different result. Indeed

$$\begin{aligned} \int_{-\infty}^{\infty} (p_n(x)p_{n-1}(x)|x|^\rho)'\, w_0(x)\, dx &= -\int_{-\infty}^{\infty} p_n(x)p_{n-1}(x)|x|^\rho w_0'(x)\, dx \\ &= 4\int_{-\infty}^{\infty} x^3 p_n(x)p_{n-1}(x)|x|^\rho w_0(x)\, dx. \end{aligned} \tag{2.10}$$

This integral can be computed by applying the three term recurrence (1.2), with $b_n = 0$, repeatedly. Indeed

$$\begin{aligned} x^2 p_n(x) &= a_{n+1} x p_{n+1}(x) + a_n x p_{n-1}(x) \\ &= a_{n+1}a_{n+2}p_{n+2}(x) + (a_{n+1}^2 + a_n^2)p_n(x) + a_n a_{n-1} p_{n-2}(x), \end{aligned}$$

and

$$\begin{aligned} x^3 p_n(x) &= a_{n+1}a_{n+2} x p_{n+2}(x) + (a_{n+1}^2 + a_n^2) x p_n(x) + a_n a_{n-1} x p_{n-2}(x) \\ &= a_{n+1}a_{n+2}a_{n+3}p_{n+3}(x) + a_{n+1}(a_{n+2}^2 + a_{n+1}^2 + a_n^2)p_{n+1}(x) \\ &\quad + a_n(a_{n+1}^2 + a_n^2 + a_{n-1}^2)p_{n-1}(x) + a_n a_{n-1} a_{n-2} p_{n-3}(x). \end{aligned}$$

From this one easily finds

$$\int_{-\infty}^{\infty} x^3 p_n(x)p_{n-1}(x)|x|^\rho w_0(x)\, dx = a_n(a_{n+1}^2 + a_n^2 + a_{n-1}^2). \tag{2.11}$$

This result holds for $n \geq 1$ if we define $a_0 = 0$. Observe that one can obtain this also by using the calculus of the Jacobi operator, since

$$(J^3)_{m,n} = \int_{-\infty}^{\infty} x^3 p_m(x)p_n(x)w_\rho(x)\, dx,$$

and the quantity of interest is $(J^3)_{n,n-1}$. This computation is quite simple since it amounts to some simple matrix multiplications. Combining (2.6) and (2.10)–(2.11) then gives

$$4a_n^2(a_{n+1}^2 + a_n^2 + a_{n-1}^2) = n + \rho\Delta_n. \tag{2.12}$$

This time we do not get a_n explicitly, but instead we get a second order non-linear recurrence relation for the recurrence coefficients. The initial conditions are $a_0 = 0$ and for a_1 we require that $p_1(x) = xp_0(x)/a_1$ has norm one, which means

$$\frac{\gamma_0^2}{a_1^2}\int_{-\infty}^{\infty} x^2 w_\rho(x)\, dx = 1,$$

and together with (1.3) this means

$$a_1 = \left(\frac{\Gamma(\frac{3+\rho}{4})}{\Gamma(\frac{1+\rho}{4})} \right)^{1/2}.$$

If we put $x_n = 2a_n^2$ then clearly $x_0 = 0$, $x_n > 0$ for $n > 0$ and

$$x_n(x_{n+1} + x_n + x_{n-1}) = n + \rho\Delta_n. \tag{2.13}$$

This recurrence relation is the *discrete Painlevé equation* d-$\mathrm{P_I}$

$$x_{n+1} + x_n + x_{n-1} = \frac{z_n + \gamma(-1)^n}{x_n} + \delta, \qquad z_n = \alpha n + \beta,$$

with $\alpha = 1$, $\beta = \rho/2$, $\gamma = -\rho/2$ and $\delta = 0$, since we can write $\Delta_n = (1 - (-1)^n)/2$.

The observation that Freud's equation (2.12) is a discrete Painlevé equation was not known to Freud but was pointed out much later by Magnus in [17]. This means that the equation has the discrete Painlevé property, which is known as *singularity confinement* (see, e.g., [8]):

Definition 2.1 (discrete Painlevé property). *If x_n is such that it results in a singularity for x_{n+1}, then there exists a $p \in \mathbb{N}$ such that this singularity is confined to $x_{n+1}, \ldots, x_{n+p}$. Furthermore x_{n+p+1} depends only on $x_{n-1}, x_{n-2}, \ldots$.*

The usual Painlevé property for differential equations is that the only movable singularities (singularities which depend on the initial conditions) of solutions of a Painlevé equation are poles. Poles are isolated singularities, hence a discrete version of poles as singularities is to require that singularities of a discrete equation are confined. This is the case for discrete Painlevé I. Consider for instance d-$\mathrm{P_I}$ in the form

$$x_n(x_{n+1} + x_n + x_{n-1}) = n,$$

then

$$x_{n+1} = \frac{n}{x_n} - x_n - x_{n-1},$$

and if $x_n = 0$ then we have a singularity for x_{n+1} which becomes $\pm\infty$. For x_{n+2} we then find $\mp\infty$ and for x_{n+3} we have the indeterminate form $(\pm\infty) + (\mp\infty)$. A more careful analysis is to put $x_n = \epsilon$ and to expand x_{n+k} in a Laurent series in ϵ. This gives

$$\begin{aligned}
x_n &= \epsilon \\
x_{n+1} &= \frac{n}{\epsilon} - x_{n-1} - \epsilon \\
x_{n+2} &= -\frac{n}{\epsilon} + x_{n-1} + \frac{n+1}{n}\epsilon + \mathcal{O}(\epsilon^2) \\
x_{n+3} &= -\frac{n+3}{n}\epsilon + \mathcal{O}(\epsilon^2) \\
x_{n+4} &= \frac{n}{n+3}x_{n-1} + \mathcal{O}(\epsilon).
\end{aligned}$$

So we see that as $\epsilon \to 0$ the indeterminate form for x_{n+3} becomes 0, but it does not give a new singularity for x_{n+4}. The singularity is confined to $x_{n+1}, x_{n+2}, x_{n+3}$.

The solution of d-P$_{\mathrm{I}}$ can not be obtained in a closed form, but one can say a few things about the behavior of the solution. Freud obtained the asymptotic behavior of the solution of (2.13) in the following way. Since $x_n \geq 0$ for $n \geq 0$ we have

$$x_n^2 \leq x_n(x_{n+1} + x_n + x_{n-1}) = n + \rho\Delta_n,$$

so that $x_n/\sqrt{n}$ is a bounded and positive sequence. Define A and B as the smallest and largest accumulation points

$$0 \leq A = \liminf_{n\to\infty} \frac{x_n}{\sqrt{n}} \leq \limsup_{n\to\infty} \frac{x_n}{\sqrt{n}} = B < \infty.$$

Choose a subsequence n' such that $x_{n'}/\sqrt{n'} \to A$, then as $n' \to \infty$ in (2.13) we have

$$1 \leq A(2B + A).$$

In a similar way we can choose a subsequence n'' such that $x_{n''}/\sqrt{n''} \to B$, and as $n' \to \infty$ in (2.13) we then find

$$B(2A + B) \leq 1.$$

Together this gives

$$B(2A + B) \leq A(2B + A),$$

so that $B^2 \leq A^2$. But since we already know that $A \leq B$, this implies that $A = B$ and hence

$$\lim_{n\to\infty} \frac{x_n}{\sqrt{n}} = A = B.$$

If we take the limit in the recurrence relation (2.13) then one finds $3A^2 = 1$ so that $A = 1/\sqrt{3}$. Recall that $x_n = 2a_n^2$, hence as a result we have

$$\lim_{n\to\infty} \frac{a_n}{n^{1/4}} = \frac{1}{\sqrt[4]{12}}. \tag{2.14}$$

The recurrence relation (2.13), with initial conditions

$$x_0 = 0, \quad x_1 = \frac{2\Gamma(\frac{3+\rho}{4})}{\Gamma(\frac{1+\rho}{4})}$$

is very unstable for computing the recurrence coefficients. Lew and Quarles [13] showed that there is a unique positive solution of the recurrence relation (2.13) with $x_0 = 0$. Hence a small error in x_1 eventually destroys the positivity of x_n. In Figure 1 we plotted the values of x_n obtained from the recurrence relation (with $\rho = 0$) by using 30 digits accuracy. The x_n are following the asymptotic behavior $\sqrt{n/3}$ quite well until $n \sim 50$ and then large deviations from the true solution appear.

Lew and Quarles proved the unicity by showing that there is an operator F acting on a Banach space of infinite sequences with $x_0 = 0$, such that the positive solution $x = (x_0, x_1, x_2, \ldots)$ of (2.13) is a fixed point: $F(x) = x$. The operator F is then shown to be a contraction, so that the fixed point is unique.

Observe that if we take a weight function of the form $w_\rho(x) = |x|^\rho \exp(-x^4 + \lambda x^2)$, then the Pearson equation becomes

$$[w_0(x)]' = (-4x^3 + 2\lambda x)w_0(x),$$



Figure 1: The result of computing x_n from d-P$_\mathrm{I}$ using 30 significant digits

and a slight modification of the previous computations gives the recurrence

$$4a_n^2(a_{n+1}^2 + a_n^2 + a_{n-1}^2) - 2\lambda a_n^2 = n + \rho\Delta_n.$$

If we put $x_n = 2a_n^2$, then the x_n satisfy the discrete Painlevé equation d-P$_\mathrm{I}$ with $\alpha = 1$, $\beta = \rho/2$, $\gamma = -\rho/2$ and $\delta = \lambda$.

2.3 Freud weight $m = 6$

For the Freud weight $w_\rho(x) = |x|^\rho \exp(-x^6)$ one can proceed in a very similar way. The Pearson equation now becomes

$$[w_0(x)]' = -6x^5 w_0(x),$$

so that

$$\int_{-\infty}^{\infty} (p_n(x)p_{n-1}(x)|x|^\rho)'\, w_0(x)\, dx = 6\int_{-\infty}^{\infty} x^5 p_n(x)p_{n-1}(x)|x|^\rho w_0(x)\, dx.$$

The integral on the right is

$$\int_{-\infty}^{\infty} x^5 p_n(x)p_{n-1}(x)|x|^\rho w_0(x)\, dx = (J^5)_{n,n-1}$$

and this is

$$\begin{aligned}(J^5)_{n,n-1} = a_n(&a_{n-2}^2 a_{n-1}^2 + a_{n-1}^4 + 2a_{n-1}^2 a_n^2 + a_{n-1}^2 a_{n+1}^2 \\ &+ a_n^4 + 2a_n^2 a_{n+1}^2 + a_{n+1}^4 + a_{n+1}^2 a_{n+2}^2).\end{aligned}$$

Together with (2.6) this gives

$$6a_n^2(a_{n-2}^2a_{n-1}^2 + a_{n-1}^4 + 2a_{n-1}^2a_n^2 + a_{n-1}^2a_{n+1}^2 \\ + a_n^4 + 2a_n^2a_{n+1}^2 + a_{n+1}^4 + a_{n+1}^2a_{n+2}^2) = n + \rho\Delta_n. \quad (2.15)$$

This is a fourth order non-linear recurrence relation for the recurrence coefficients. It is within the hierarchy of discrete Painlevé I [6]. Freud was also able to obtain the asymptotic behavior for the a_n in this case. Obviously

$$6a_n^6 \le 6a_n^2(a_{n-2}^2a_{n-1}^2+a_{n-1}^4+2a_{n-1}^2a_n^2+a_{n-1}^2a_{n+1}^2+a_n^4+2a_n^2a_{n+1}^2+a_{n+1}^4+a_{n+1}^2a_{n+2}^2) = n+\rho\Delta_n,$$

hence $a_n/n^{1/6}$ is a positive and bounded sequence. If we define

$$0 \le A = \liminf_{n\to\infty} \frac{a_n^2}{n^{1/3}} \le \limsup_{n\to\infty} \frac{a_n^2}{n^{1/3}} = B < \infty,$$

then by taking a subsequence that converges to A we find

$$6A(5B^2 + 4AB + A) \ge 1$$

and by taking a subsequence that converges to B we find

$$6B(5A^2 + 4AB + B^2) \le 1.$$

Combining both inequalities gives

$$6B(5A^2 + 4AB + B^2) \le 6A(5B^2 + 4AB + A)$$

so that $A^2B + B^3 \le AB^2 + A^3$. This is equivalent with

$$(B - A)(A^2 + AB + B^2) \le (B - A)AB.$$

If we assume that $A < B$, then this would imply that $A^2 + B^2 \le 0$, or $A = B = 0$, which is impossible (since $A < B$). Hence our assumption is false and $A = B$. Taking the limit in (2.15) gives $60A^3 = 1$, hence

$$\lim_{n\to\infty} \frac{a_n}{n^{1/6}} = \frac{1}{\sqrt[6]{60}}. \quad (2.16)$$

2.4 Freud's conjecture

On the basis of (2.9), (2.14) and (2.16), Freud made the conjecture that for every $m > 0$ and $\rho > -1$ one has

$$\lim_{n\to\infty} \frac{a_n}{n^{1/m}} = \left(\frac{\Gamma(m+1)}{\Gamma(\frac{m}{2})\Gamma(\frac{m}{2}+1)}\right)^{-1/m}. \quad (2.17)$$

Furthermore Freud showed that if the limit exists for even m, then it is equal to the expression in (2.17). Freud could not prove the existence of the limit for even $m \ge 8$ because the central coefficient a_n in the non-linear recurrence relation does not occur sufficiently often and more non central coefficients $a_{n\pm k}$ with $k \ne 0$ appear, making the recurrence no longer 'diagonally dominant'. The simple trick using lim sup and lim inf then no longer suffices to show that the limit exists. The proof of Freud's conjecture for every even integer m was given by Alphonse Magnus [15] [16]. His proof still consists of obtaining a non-linear recurrence relation for the a_n (the Freud equation, which is within the hierarchy of discrete Painlevé I), but a more subtle argument is used to prove the existence of the limit. Freud's conjecture for general $m > 0$ was finally proved by Lubinsky, Mhaskar and Saff [14]. The proof for general $m > 0$ no longer uses a recurrence relation for the recurrence coefficients but relies on the Mhaskar-Rakhmanov-Saff number and results of weighted polynomial approximation and an equilibrium problem of logarithmic potential theory with external field.

For m an even positive integer, Máté, Nevai and Zaslavsky [18] obtained an asymptotic expansion of the form

$$\frac{a_n^2}{n^{1/m}} = \sum_{k=0}^{\infty} \frac{c_k}{n^{2k}},$$

where c_0 is the constant in Freud's conjecture (2.17), but the other coefficients c_k with $k > 0$ are not explicitly known. Their analysis is again based on the non-linear recurrence relation for the recurrence coefficients.

3 Orthogonal polynomials on the unit circle

In this section we will consider orthogonal polynomials on the unit circle $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$. A very good source for the general theory is the recent set of books by Barry Simon [22]. The sequence of polynomials $\{\varphi_n, n = 0, 1, 2, \ldots\}$ is orthonormal on the unit circle with respect to a weight w if

$$\frac{1}{2\pi} \int_0^{2\pi} \varphi_n(z)\overline{\varphi_m(z)} w(\theta)\, d\theta = \delta_{n,m}. \tag{3.1}$$

These polynomials are unique if we agree to make the leading coefficient positive:

$$\varphi_n(z) = \kappa_n z^n + \cdots, \qquad \kappa_n > 0.$$

The monic polynomials are usually denoted by $\Phi_n(z) = \varphi_n(z)/\kappa_n$. An important property, which replaces the three term recurrence relation for orthogonal polynomials on the real line, is the Szegő recurrence

$$z\Phi_n(z) = \Phi_{n+1}(z) + \overline{\alpha_n}\Phi_n^*(z), \tag{3.2}$$

where $\Phi_n^*(z) = z^n \overline{\Phi}_n(1/z)$ is the reversed polynomial and $\overline{\Phi}_n$ is the polynomial Φ_n but with complex conjugated coefficients. In [22] the recurrence coefficients α_n $(n = 0, 1, 2, 3, \ldots)$ are called Verblunsky coefficients. They are given by $\alpha_n = -\overline{\Phi_{n+1}(0)}$ and they satisfy $|\alpha_n| < 1$ for $n \geq 0$ and $\alpha_{-1} = -1$. An important relation between κ_n and α_n was found by Szegő:

$$\kappa_n^2 = \sum_{k=0}^{n} |\varphi_k(0)|^2,$$

from which it follows that

$$\kappa_{n+1}^2 - \kappa_n^2 = |\varphi_{n+1}(0)|^2 = \kappa_{n+1}^2 |\alpha_n|^2,$$

so that

$$\frac{\kappa_n^2}{\kappa_{n+1}^2} = 1 - |\alpha_n|^2. \tag{3.3}$$

3.1 Modified Bessel polynomials

We will take a closer look at the orthogonal polynomials on the unit circle for the weight

$$w(\theta) = \exp(\lambda \cos\theta), \qquad \theta \in [-\pi, \pi].$$

observe that $w(-\theta) = w(\theta)$, which implies that the Verblunsky coefficients are real. Ismail [11, pp. 236–239] calls the resulting orthogonal polynomials the *modified Bessel polynomials* since the trigonometric moments of this weight are in terms of the modified Bessel function

$$\frac{1}{2\pi}\int_0^{2\pi} e^{in\theta} w(\theta)\, d\theta = \frac{1}{\pi}\int_0^{\pi} \cos n\theta \exp(\lambda\cos\theta)\, d\theta = I_n(\lambda).$$

These polynomials appear in the analysis of unitary random matrices [20, 23, 9] and play an important role in the asymptotic distribution of the length of the longest increasing subsequence of random permutations [2].

Periwal and Shevitz [20] found a non-linear recurrence relation for the Verblunsky coefficients of these orthogonal polynomials (see also [9, 23]). If $z = e^{i\theta}$ then

$$w(\theta) = \exp\left(\frac{\lambda}{2}(z + \frac{1}{z})\right) := \hat{w}(z),$$

and this weight satisfies a Pearson equation of the form

$$\hat{w}'(z) = \frac{\lambda}{2}(1 - \frac{1}{z^2})\hat{w}(z). \tag{3.4}$$

Consider the integral

$$\frac{1}{2\pi i}\int_{\mathbb{T}} \overline{\varphi_n(z)}\varphi_{n+1}(z)\hat{w}'(z)\, \frac{dz}{z}, \tag{3.5}$$

then by means of Pearson's equation (3.4) we find

$$\begin{aligned} \frac{1}{2\pi i}\int_{\mathbb{T}} \overline{\varphi_n(z)}\varphi_{n+1}(z)\hat{w}'(z)\, \frac{dz}{z} &= \frac{\lambda}{4\pi i}\int_{\mathbb{T}} \overline{\varphi_n(z)}\varphi_{n+1}(z)(1 - \frac{1}{z^2})\hat{w}(z)\, \frac{dz}{z} \\ &= \frac{\lambda}{4\pi}\int_0^{2\pi} \overline{\varphi_n(z)}\varphi_{n+1}(z) w(\theta)\, d\theta \\ &\quad - \frac{\lambda}{4\pi}\int_0^{2\pi} \overline{\varphi_n(z) z^2}\varphi_{n+1}(z) w(\theta)\, d\theta. \end{aligned}$$

The first integral on the right is zero because of orthogonality. For the second integral we use the recurrence (3.2) for the orthonormal polynomials

$$z\varphi_n(z) = \frac{\kappa_n}{\kappa_{n+1}}\varphi_{n+1}(z) + \overline{\alpha_n}\varphi_n^*(z) \tag{3.6}$$

to find

$$\begin{aligned}\frac{\lambda}{4\pi}\int_0^{2\pi}\overline{z^2\varphi_n(z)}\varphi_{n+1}(z)w(\theta)\,d\theta &= \frac{\kappa_n}{\kappa_{n+1}}\frac{\lambda}{4\pi}\int_0^{2\pi}\overline{z\varphi_{n+1}(z)}\varphi_{n+1}(z)w(\theta)\,d\theta \\ &\quad + \alpha_n\frac{\lambda}{4\pi}\int_0^{2\pi}\overline{z\varphi_n^*(z)}\varphi_{n+1}(z)w(\theta)\,d\theta.\end{aligned}$$

If we use (3.6) for $n+1$ and orthogonality, then

$$\begin{aligned}\frac{\kappa_n}{\kappa_{n+1}}\frac{\lambda}{4\pi}\int_0^{2\pi}\overline{z\varphi_{n+1}(z)}\varphi_{n+1}(z)w(\theta)\,d\theta &= \alpha_{n+1}\frac{\kappa_n}{\kappa_{n+1}}\frac{\lambda}{4\pi}\int_0^{2\pi}\overline{\varphi_{n+1}^*(z)}\varphi_{n+1}(z)w(\theta)\,d\theta \\ &= -\frac{\lambda}{2}\frac{\kappa_n}{\kappa_{n+1}}\alpha_{n+1}\overline{\alpha_n}\end{aligned}$$

because

$$\begin{aligned}\varphi_{n+1}^*(z) &= \frac{\overline{\varphi_{n+1}(0)}}{\kappa_{n+1}}\varphi_{n+1}(z) + \text{ lower degree terms} \\ &= -\alpha_n\varphi_{n+1}(z) + \text{ lower degree terms.}\end{aligned}$$

In a similar way we have

$$z\varphi_n^*(z) = -\alpha_{n-1}\frac{\kappa_n}{\kappa_{n+1}}\varphi_{n+1}(z) + \text{ lower degree terms}$$

so that

$$\alpha_n\frac{\lambda}{4\pi}\int_0^{2\pi}\overline{z\varphi_n^*(z)}\varphi_{n+1}(z)w(\theta)\,d\theta = -\frac{\lambda}{2}\alpha_n\overline{\alpha_{n-1}}\frac{\kappa_n}{\kappa_{n+1}}.$$

Combining all these results gives

$$\frac{1}{2\pi i}\int_{\mathbb{T}}\overline{\varphi_n(z)}\varphi_{n+1}(z)\hat{w}'(z)\,\frac{dz}{z} = \frac{\lambda}{2}\frac{\kappa_n}{\kappa_{n+1}}(\alpha_{n+1}\overline{\alpha_n} + \alpha_n\overline{\alpha_{n-1}}). \tag{3.7}$$

We can compute this integral also using integration by parts, to find

$$\frac{1}{2\pi i}\int_{\mathbb{T}}\overline{\varphi_n(z)}\varphi_{n+1}(z)\hat{w}'(z)\,\frac{dz}{z} = -\frac{1}{2\pi i}\int_{\mathbb{T}}[\overline{z\varphi_n(z)}\varphi_{n+1}(z)]'\hat{w}(z)\,dz.$$

We have to be a little bit careful because $\overline{z\varphi_n(z)}$ is not analytic in the complex plane, but on the unit circle $\mathbb{T}$ we have $\overline{\varphi_n(z)} = z^{-n}\varphi_n^*(z)$ so that

$$\begin{aligned}\frac{1}{2\pi i}\int_{\mathbb{T}}[\overline{z\varphi_n(z)}\varphi_{n+1}(z)]'\hat{w}(z)\,dz &= \frac{1}{2\pi i}\int_{\mathbb{T}}[z^{-n-1}\varphi_n^*(z)\varphi_{n+1}(z)]'\hat{w}(z)\,dz \\ &= -\frac{n+1}{2\pi i}\int_{\mathbb{T}}z^{-n-1}\varphi_n^*(z)\varphi_{n+1}(z)\hat{w}(z)\,\frac{dz}{z} \\ &\quad + \frac{1}{2\pi i}\int_{\mathbb{T}}z^{-n}[\varphi_n^*]'(z)\varphi_{n+1}(z)\hat{w}(z)\,\frac{dz}{z} \\ &\quad + \frac{1}{2\pi i}\int_{\mathbb{T}}z^{-n}\varphi_n^*(z)\varphi_{n+1}'(z)\hat{w}(z)\,\frac{dz}{z}.\end{aligned}$$

If we use the recurrence relation (3.6) then

$$\begin{aligned}\frac{1}{2\pi i}\int_{\mathbb{T}} z^{-n-1}\varphi_n^*(z)\varphi_{n+1}(z)\hat{w}(z)\,\frac{dz}{z} &= \frac{1}{2\pi}\int_0^{2\pi}\overline{z\varphi_n(z)}\varphi_{n+1}(z)w(\theta)\,d\theta \\ &= \frac{\kappa_n}{\kappa_{n+1}},\end{aligned}$$

by orthogonality we find

$$\begin{aligned}\frac{1}{2\pi i}\int_{\mathbb{T}} z^{-n}[\varphi_n^*]'(z)\varphi_{n+1}(z)\hat{w}(z)\,\frac{dz}{z} &= \frac{1}{2\pi}\int_0^{2\pi}\overline{z\varphi_n'(z)}\varphi_{n+1}(z)w(\theta)\,d\theta \\ &= 0,\end{aligned}$$

and if we use

$$\varphi_{n+1}'(z) = (n+1)\frac{\kappa_{n+1}}{\kappa_n}\varphi_n(z) + \text{ lower degree terms,}$$

then

$$\begin{aligned}\frac{1}{2\pi i}\int_{\mathbb{T}} z^{-n}\varphi_n^*(z)\varphi_{n+1}'(z)\hat{w}(z)\,\frac{dz}{z} &= \frac{1}{2\pi}\int_0^{2\pi}\overline{\varphi_n(z)}\varphi_{n+1}'(z)w(\theta)\,d\theta \\ &= (n+1)\frac{\kappa_{n+1}}{\kappa_n}.\end{aligned}$$

These computations give

$$\frac{1}{2\pi i}\int_{\mathbb{T}}[\overline{z\varphi_n(z)}\varphi_{n+1}(z)]'\hat{w}(z)\,dz = (n+1)\frac{\kappa_{n+1}}{\kappa_n}\left(1-\frac{\kappa_n^2}{\kappa_{n+1}^2}\right). \tag{3.8}$$

Now we can combine (3.7) and (3.8) to find

$$-\frac{\lambda}{2}\frac{\kappa_n}{\kappa_{n+1}}(\alpha_{n+1}\overline{\alpha_n}+\alpha_n\overline{\alpha_{n-1}}) = (n+1)\frac{\kappa_{n+1}}{\kappa_n}\left(1-\frac{\kappa_n^2}{\kappa_{n+1}^2}\right)$$

which, together with (3.3) gives

$$-\frac{\lambda}{2}(1-|\alpha_n|^2)(\alpha_{n+1}\overline{\alpha_n}+\alpha_n\overline{\alpha_{n-1}}) = (n+1)|\alpha_n|^2.$$

Recall that $w(-\theta)=w(\theta)$ implies that the α_n are real. Hence when $\alpha_n \neq 0$ then

$$-\frac{\lambda}{2}(1-\alpha_n^2)(\alpha_{n+1}+\alpha_{n-1}) = (n+1)\alpha_n. \tag{3.9}$$

This non-linear recurrence relation corresponds to the discrete Painlevé equation d-P$_{\text{II}}$

$$x_{n+1}+x_{n-1} = \frac{x_n(\alpha n+\beta)+\gamma}{1-x_n^2}$$

with $\alpha_n = x_n$, $\alpha=\beta=-2/\lambda$ and $\gamma=0$. The initial values are

$$\alpha_{-1}=-1, \qquad \alpha_0 = \frac{\int_0^{2\pi} z w(\theta)\,d\theta}{\int_0^{2\pi} w(\theta)\,d\theta} = \frac{I_1(\lambda)}{I_0(\lambda)}.$$

4 Discrete orthogonal polynomials

In this section we will study certain discrete orthogonal polynomials on the integers $\mathbb{N}$. The orthonormality now becomes

$$\sum_{k=0}^{\infty} p_n(k)p_m(k)w_k = \delta_{n,m}, \qquad n,m \geq 0. \tag{4.1}$$

Instead of the differential operator we will now be using difference operators, namely the forward difference Δ and the backward difference ∇ for which

$$\Delta f(x) = f(x+1) - f(x), \qquad \nabla f(x) = f(x) - f(x-1).$$

We now have two sequences $\{a_n : n = 1,2,\ldots\}$ and $\{b_n : n = 0,1,2,\ldots\}$ of recurrence coefficients, and we need two recurrence relations to determine all a_n and b_n.

4.1 Charlier polynomials

Charlier polynomials are the orthonormal polynomials for the Poisson distribution

$$w_k = \frac{a^k}{k!}, \qquad k \in \mathbb{N},\ a > 0.$$

Observe that

$$w_{k-1} = \frac{k}{a}\, w_k \tag{4.2}$$

which is the (discrete) Pearson equation for the Poisson distribution. It can also be written as $a\nabla w_k = (a-k)w_k$. The Pearson equation gives the following structure relation for Charlier polynomials.

Lemma 4.1. *For the orthonormal Charlier polynomials one has*

$$p_n(x+1) = p_n(x) + \frac{a_n}{a} p_{n-1}(x), \tag{4.3}$$

where a_n are the coefficients in the recurrence relation (1.2).

Proof. If we expand $p_n(x+1)$ into a Fourier series, then

$$p_n(x+1) = \sum_{j=0}^{n} A_{n,j} p_j(x),$$

and if we compare the leading coefficients then $A_{n,n} = 1$. The other Fourier coefficients are given by

$$A_{n,j} = \sum_{k=0}^{\infty} p_n(k+1)p_j(k)w_k = \sum_{k=1}^{\infty} p_n(k)p_j(k-1)w_{k-1},$$

and if we use (4.2) then this gives

$$A_{n,j} = \frac{1}{a}\sum_{k=0}^{\infty} k p_n(k)p_j(k-1)w_k.$$

The polynomial $xp_j(x-1)$ has degree $j+1$, hence by orthogonality $A_{n,j} = 0$ whenever $j < n-1$. For $j = n-1$ we have

$$xp_{n-1}(x-1) = \frac{\gamma_{n-1}}{\gamma_n} p_n(x) + \text{ lower degree terms}$$

so that (1.4) gives the desired result. $\square$

Note that we can write (4.3) also as

$$\Delta p_n(x) = \frac{a_n}{a} p_{n-1}(x).$$

If we compare the leading coefficient in the latter, then $n\gamma_n = a_n\gamma_{n-1}/a$, so that

$$a_n^2 = an.$$

We will now compute the sum

$$\sum_{k=0}^{\infty} p_n^2(k+1) w_k$$

in two different ways. If we use the Pearson equation (4.2) then

$$\sum_{k=0}^{\infty} p_n^2(k+1) w_k = \sum_{k=1}^{\infty} p_n^2(k) w_{k-1} = \frac{1}{a} \sum_{k=0}^{\infty} k p_n^2(k) w_k = \frac{b_n}{a}.$$

On the other hand, the structure relation (4.3) gives

$$\sum_{k=0}^{\infty} p_n^2(k+1) w_k = \sum_{k=0}^{\infty} \Big(p_n(k) + \frac{a_n}{a} p_{n-1}(k) \Big)^2 w_k = 1 + \frac{a_n^2}{a^2}.$$

Combining both computations gives

$$b_n = a + \frac{a_n^2}{a} = n + a.$$

These simple computations show that the recurrence coefficients for Charlier polynomials are given by

$$a_n = \sqrt{an}, \qquad b_n = n + a.$$

4.2 Generalized Charlier polynomials

If we take the weights

$$w_k = \frac{a^k}{(k!)^N}, \qquad k \in \mathbb{N},\ a > 0,$$

with $N \in \{1, 2, 3, \ldots\}$, then for $N = 1$ one finds the Charlier polynomials and for $N \geq 2$ the generalized Charlier polynomials. These were introduced by Hounkonnou et al. in [10]. The Pearson equation is

$$w_{k-1} = \frac{k^N}{a}\, w_k, \tag{4.4}$$

which can also be written as $a\nabla w_k = (a - k^N) w_k$. For $N \geq 2$ the factor $a - k^N$ is a polynomial in k of degree greater than one, and hence the weight is no longer classical but semi-classical. We will investigate the case $N = 2$ in more detail.

Lemma 4.2. *For $N = 2$ the generalized Charlier polynomials satisfy the structure relation*

$$p_n(x+1) = p_n(x) + \frac{n}{a_n} p_{n-1}(x) + \frac{a_n a_{n-1}}{a} p_{n-2}(x), \tag{4.5}$$

where a_n are the recurrence coefficients in the three-term recurrence relation (1.2).

Proof. If we expand $p_n(x+1)$ into a Fourier series, then

$$p_n(x+1) = \sum_{j=0}^{n} A_{n,j} p_j(x).$$

Comparing coefficients of x^n gives $A_{n,n} = 1$, and comparing coefficients of x^{n-1} gives $A_{n,n-1} = n/a_n$. The remaining Fourier coefficients are given by

$$A_{n,j} = \sum_{k=0}^{\infty} p_n(k+1) p_j(k) w_k = \sum_{k=1}^{\infty} p_n(k) p_j(k-1) w_{k-1}.$$

If we use the Pearson equation (4.4) then

$$A_{n,j} = \frac{1}{a} \sum_{k=0}^{\infty} p_n(k) p_j(k-1) k^2 w_k.$$

The polynomial $x^2 p_j(x-1)$ is of degree $j+2$ and hence by orthogonality $A_{n,j} = 0$ for $j < n-2$. For $j = n-2$ we have

$$x^2 p_{n-2}(x-1) = \frac{\gamma_{n-2}}{\gamma_n} p_n(x) + \text{ lower degree terms}$$

so that

$$A_{n,n-2} = \frac{1}{a} \frac{\gamma_{n-2}}{\gamma_n} = \frac{a_n a_{n-1}}{a},$$

where we used (1.4), which gives the required result. □

The structure relation (4.5) can also be written as

$$\Delta p_n(x) = \frac{n}{a_n} p_{n-1}(x) + \frac{a_n a_{n-1}}{a} p_{n-2}(x).$$

If we compare coefficients of x^{n-2}, where we use

$$p_n(x) = \gamma_n x^n + \delta_n x^{n-1} + \cdots ,$$

then we find

$$\binom{n}{2} \gamma_n + (n-1)\delta_n = \frac{n}{a_n} \delta_{n-1} + \frac{a_n a_{n-1}}{a} \gamma_{n-2}. \tag{4.6}$$

If $x_{1,n} < x_{2,n} < \cdots < x_{n,n}$ are the zeros of p_n, then by Viète's symmetric formulas we have

$$\frac{\delta_n}{\gamma_n} = -\sum_{k=1}^{n} x_{k,n}.$$

The zeros of p_n are equal to the eigenvalues of the truncated Jacobi matrix

$$J_n = \begin{pmatrix} b_0 & a_1 & & & & \\ a_1 & b_1 & a_2 & & & \\ & a_2 & b_2 & a_3 & & \\ & & a_3 & b_3 & \ddots & \\ & & & \ddots & \ddots & a_{n-1} \\ & & & & a_{n-1} & b_{n-1} \end{pmatrix}$$

and the sum of all eigenvalues is the trace of the matrix, hence

$$\frac{\delta_n}{\gamma_n} = -\sum_{j=0}^{n-1} b_j.$$

If we use this in (4.6), then

$$\binom{n}{2} - nb_{n-1} + \sum_{j=0}^{n-1} b_j = \frac{a_n^2 a_{n-1}^2}{a}. \tag{4.7}$$

In order to get rid of the non-homogeneous terms, we put $b_n = n + \hat{b}_n$, and the relation becomes

$$-n\hat{b}_{n-1} + \sum_{j=0}^{n-1} \hat{b}_j = \frac{a_n^2 a_{n-1}^2}{a}.$$

Differencing both sides gives

$$-na(\hat{b}_n - \hat{b}_{n-1}) = a_n^2(a_{n+1}^2 - a_{n-1}^2), \tag{4.8}$$

which may be considered as the first Freud equation for the recurrence coefficients.

Next, we will compute

$$\sum_{k=0}^{\infty} p_n(k+1)p_{n-1}(k+1)w_k$$

in two different ways. First we use the Pearson equation (4.4) to find

$$\begin{aligned} \sum_{k=0}^{\infty} p_n(k+1)p_{n-1}(k+1)w_k &= \sum_{k=1}^{\infty} p_n(k)p_{n-1}(k)w_{k-1} \\ &= \frac{1}{a}\sum_{k=0}^{\infty} k^2 p_n(k)p_{n-1}(k)w_k \\ &= \frac{1}{a}(J^2)_{n,n-1}. \end{aligned}$$

The entry $(J^2)_{n,n-1}$ can be computed easily by repeatedly using the recurrence relation (1.2) and is equal to $(J)^2_{n,n-1} = a_n(b_n + b_{n-1})$, so that

$$\sum_{k=0}^{\infty} p_n(k+1)p_{n-1}(k+1)w_k = \frac{a_n(b_n + b_{n-1})}{a}. \tag{4.9}$$

On the other hand, we can use the structure relation (4.5) to find

$$\begin{aligned}\sum_{k=0}^{\infty} p_n(k+1)p_{n-1}(k+1)w_k &= \sum_{k=0}^{\infty}\left(p_n(k)+\frac{n}{a_n}p_{n-1}(k)+\frac{a_na_{n-1}}{a}p_{n-2}(k)\right)\\ &\quad\times\left(p_{n-1}(k)+\frac{n-1}{a_{n-1}}p_{n-2}(k)+\frac{a_{n-1}a_{n-2}}{a}p_{n-3}(k)\right)w_k\\ &= \frac{n}{a_n}+\frac{(n-1)a_n}{a}, \qquad (4.10)\end{aligned}$$

where the last equality follows from the orthonormality (4.1). Combining (4.9) and (4.10), and recalling that $b_n = n + \hat{b}_n$, then gives the second Freud equation

$$a_n^2(\hat{b}_n + \hat{b}_{n-1} + n) = na. \qquad (4.11)$$

If we eliminate na from the two equation (4.8) and (4.11), then

$$-(\hat{b}_n + \hat{b}_{n-1} + n)(\hat{b}_n - \hat{b}_{n-1}) = a_{n+1}^2 - a_{n-1}^2.$$

Summing both sides of this equation gives

$$-\hat{b}_n^2 + \sum_{k=0}^{n-1}\hat{b}_k - n\hat{b}_n = a_{n+1}^2 + a_n^2 - a. \qquad (4.12)$$

Summing both sides of (4.8) gives

$$a\left(\sum_{k=0}^{n-1}\hat{b}_k - n\hat{b}_n\right) = a_n^2a_{n+1}^2. \qquad (4.13)$$

Combining (4.12) and (4.13) then gives

$$a_n^2a_{n+1}^2 - a(a_{n+1}^2 + a_n^2) + a^2 = a\hat{b}_n^2,$$

which is equivalent to

$$a\hat{b}_n^2 = (a_{n+1}^2 - a)(a_n^2 - a). \qquad (4.14)$$

This means that $a_n^2 - a$ and $a_{n+1}^2 - a$ have the same sign, and since $a_0 = 0$ we must conclude that $a_n^2 - a < 0$ for $n \geq 0$. We may therefore write

$$a_n^2 = a(1 - c_n^2), \qquad (4.15)$$

with $c_0 = 1$, and then (4.14) becomes

$$\hat{b}_n = \sqrt{a}c_nc_{n+1}. \qquad (4.16)$$

The second Freud equation (4.11) becomes

$$(1 - c_n^2)\sqrt{a}(c_{n+1} + c_{n-1}) = nc_n. \qquad (4.17)$$

If we compute the coefficients c_n from the recurrence relation (4.17), then we obtain the recurrence coefficients $b_n = n + \hat{b}_n$ from (4.16) and the a_n from (4.15). The non-linear recurrence relation (4.17) corresponds to the discrete Painlevé II equation

$$x_{n+1} + x_{n-1} = \frac{x_n(\alpha n + \beta) + \gamma}{1 - x_n^2}$$

with $c_n = x_n$ and $\alpha = 1/\sqrt{a}$, $\beta = \gamma = 0$. We need to find the solution with $c_0 = 1$ and $c_1^2 = 1 - a_1^2/a$. Observe that if we require that $p_1(x) = (x - b_0)p_0/a_1$ is orthogonal to p_0, then

$$\sum_{k=0}^{\infty} (k - b_0) w_k = 0$$

so that

$$b_0 = \frac{\sum_{k=0}^{\infty} k \frac{a^k}{(k!)^2}}{\sum_{k=0}^{\infty} \frac{a^k}{(k!)^2}} = \frac{\sqrt{a} I_1(2\sqrt{a})}{I_0(2\sqrt{a})},$$

where

$$I_\nu(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{2k+\nu}}{k!\Gamma(k+\nu+1)}$$

is the modified Bessel function. From (4.16) we then see that

$$c_1 = \frac{I_1(2\sqrt{a})}{I_0(2\sqrt{a})}. \tag{4.18}$$

The non-linear recurrence relation (4.17) with initial conditions $c_0 = 1$ and $c_1 = I_1(2\sqrt{a})/I_0(2\sqrt{a})$ is again very unstable for computing all the c_n recursively. One can show that the discrete Painlevé equation with $\gamma = 0$ and $c_0 = \pm 1$ has only one solution for which $-1 < c_n < 1$ for all $n \geq 1$ (see [24]), and this is the solution that we need since $a_n^2 = a(1 - c_n^2)$ needs to be positive. Hence a slight deviation from the actual initial value c_1 will destroy the positivity of the a_n^2 eventually. In Figure 2 we have plotted the c_n obtained from the recurrence relation with an accuracy of 30 digits. The c_n converge quickly to zero, but for n near 40 we see that the c_n deviate quite a lot from zero.

The discrete Painlevé II equation also arose in Section 3 for the Verblunsky coefficients of certain orthogonal polynomials on the unit circle. Verblunsky coefficients always have the property that $|\alpha_n| < 1$ for $n \geq 0$, hence in that case one also requires the unique solution with $\alpha_{-1} = -1$ for which $-1 < \alpha_n < 1$ for $n \geq 0$. Observe that there is a shift in the index since we are using Verblunsky coefficients, in which case the recurrence starts with $\alpha_{-1} = -1$.

Obviously the equation (4.17) satisfies the discrete Painlevé property. Indeed, we have

$$c_{n+1} = \frac{n c_n/\sqrt{a}}{1 - c_n^2} - c_{n-1},$$

hence a singularity will appear in c_{n+1} whenever $c_n = \pm 1$. A careful analysis gives that for c_n near 1

$$\begin{aligned}
c_n &= 1 + \epsilon \\
c_{n+1} &= -\frac{n}{2\sqrt{a}} \frac{1}{\epsilon} - \frac{n}{4\sqrt{a}} - c_{n-1} + \mathcal{O}(\epsilon) \\
c_{n+2} &= -1 + \frac{n+2}{n} \epsilon + \mathcal{O}(\epsilon^2) \\
c_{n+3} &= \frac{n+1}{\sqrt{a}(n+2)} - \frac{n}{n+2} c_{n-1} + \mathcal{O}(\epsilon),
\end{aligned}$$



Figure 2: The result of computing c_n from d-$\mathrm{P_{II}}$ using 30 significant digits

and near -1

$$\begin{aligned} c_n &= -1+\epsilon \\ c_{n+1} &= -\frac{n}{2\sqrt{a}}\,\frac{1}{\epsilon}+\frac{n}{4\sqrt{a}}-c_{n-1}+\mathcal{O}(\epsilon) \\ c_{n+2} &= 1+\frac{n+2}{n}\,\epsilon+\mathcal{O}(\epsilon^2) \\ c_{n+3} &= -\frac{n+1}{\sqrt{a}(n+2)}-\frac{n}{n+2}c_{n-1}+\mathcal{O}(\epsilon), \end{aligned}$$

so that in both cases the singularity is confined to c_{n+1} and c_{n+2}. Observe that the critical value 1 for c_n results in the critical value -1 for c_{n+2}, and that the critical value -1 for c_n results in the critical value 1 for c_{n+2}.

5 q-Orthogonal polynomials

Here we consider orthogonal polynomials on the exponential lattice $\{\pm q^n, n\in\mathbb{N}\}$, where $0<q<1$. The orthogonality is of the form

$$\int_{-1}^{1} p_n(x)p_m(x)w(x)\,d_qx=\delta_{m,n}, \tag{5.1}$$

where the q-integral is defined by

$$\int_{-1}^{1} f(x)\,d_qx=(1-q)\sum_{k=0}^{\infty} f(q^k)q^k+(1-q)\sum_{k=0}^{\infty} f(-q^k)q^k.$$

We will only consider even weights for which $w(-x) = w(x)$, in which case the orthogonal polynomials have the symmetry property $p_n(-x) = (-1)^n p_n(x)$, i.e., the polynomials are even when n is even and odd when n is odd. The recurrence relation will then be of the form

$$xp_n(x) = a_{n+1}p_{n+1}(x) + a_n p_{n-1}(x), \tag{5.2}$$

with $p_{-1} = 0$. The results in this section were obtained for the first time by Nijhoff [19], but we take a slightly different approach.

5.1 Discrete q-Hermite I polynomials

The orthonormal discrete q-Hermite I polynomials [12, §3.28] are given by

$$\int_{-1}^{1} p_n(x)p_m(x)(qx;q)_\infty(-qx;q)_\infty \, d_qx = \delta_{m,n},$$

where

$$(x;q)_\infty = \prod_{k=0}^{\infty}(1 - xq^k).$$

Observe that

$$(qx;q)_\infty(-qx;q)_\infty = \prod_{k=0}^{\infty}(1 - x^2q^{2k}) = (x^2q^2;q^2)_\infty,$$

so that the weight can be defined as $w(x) = (x^2q^2;q^2)_\infty$. In terms of the q-exponential function $E_q(z) = (-z;q)_\infty$ we have $w(x) = E_{q^2}(-x^2q^2)$, and since $E_q((1-q)z) \to \exp(z)$ when $q \to 1$ it follows that $w(\sqrt{1-q^2}x) \to \exp(-x^2)$ when $q \to 1$, which shows that this weight is a q-analog of the Hermite weight. One easily finds that

$$(1 - x^2)w(x) = w(x/q), \tag{5.3}$$

which is the Pearson equation for this weight on the q-lattice. The structure relation for the corresponding orthogonal polynomials is in terms of the q-difference operator D_q for which

$$D_qf(x) = \begin{cases} \dfrac{f(qx) - f(x)}{x(q-1)}, & \text{if } x \neq 0, \\ f'(0), & \text{if } x = 0. \end{cases}$$

Lemma 5.1. *The discrete q-Hermite I polynomials satisfy*

$$D_qp_n(x) = \frac{a_n}{q^{n-1}(1-q)} \, p_{n-1}. \tag{5.4}$$

Proof. Clearly $D_qp_n(x)$ is a polynomial of degree $n-1$ and $D_qp_n(-x) = (-1)^{n-1}D_qp_n(x)$. If we expand this polynomial into a Fourier series, then

$$D_qp_n(x) = \sum_{j=0}^{n-1} a_{j,n}p_j(x),$$

with

$$a_{j,n} = \int_{-1}^{1} D_qp_n(x) \, p_j(x)w(x) \, d_qx.$$

The symmetry shows that $a_{j,n} = 0$ whenever $n - j$ is even. When $n - j$ is odd then

$$\begin{aligned} a_{j,n} &= 2(1-q) \sum_{k=0}^{\infty} D_q p_n(q^k)\ p_j(q^k) q^k w(q^k) \\ &= -2 \sum_{k=0}^{\infty} [p_n(q^{k+1}) - p_n(q^k)] p_j(q^k) w(q^k) \\ &= -2 \sum_{k=0}^{\infty} p_n(q^{k+1}) p_j(q^k) w(q^k) + 2 \sum_{k=0}^{\infty} p_n(q^k) p_j(q^k) w(q^k). \end{aligned}$$

Both sums are finite since either p_n or p_j is an odd polynomial. Using the Pearson equation (5.3), and a shift in the summation index in the first sum, gives

$$\begin{aligned} a_{j,n} &= -2 \sum_{k=0}^{\infty} p_n(q^k) p_j(q^{k-1}) w(q^k)(1 - q^{2k}) + 2 \sum_{k=0}^{\infty} p_n(q^k) p_j(q^k) w(q^k) \\ &= \int_{-1}^{1} p_n(x) \frac{p_j(x/q) - p_j(x)}{x(q-1)} w(x)\, d_q x + \frac{1}{1-q} \int_{-1}^{1} x p_n(x) p_j(x/q) w(x)\, d_q x. \end{aligned}$$

The first integral on the right is zero because of orthogonality. The second integral only gives a contribution when $j = n - 1$, in which case

$$a_{n-1,n} = \frac{1}{1-q} \int_{-1}^{1} x p_n(x) p_{n-1}(x/q) w(x)\, d_q x.$$

The recurrence relation (5.2) gives

$$x p_{n-1}(x/q) = q a_n p_n(x/q) + q a_{n-1} p_{n-2}(x/q),$$

and since

$$p_n(x/q) = q^{-n} p_n(x) + \text{ lower degree terms}$$

this gives

$$a_{n-1,n} = \frac{q a_n}{1-q} \int_{-1}^{1} p_n(x) p_n(x/q) w(x)\, d_q x = \frac{a_n}{q^{n-1}(1-q)},$$

which gives the desired structure relation. □

If we compare the leading coefficients on both sides of (5.4), then

$$\gamma_n \frac{1-q^n}{1-q} = \gamma_{n-1} \frac{a_n}{q^{n-1}(1-q)},$$

so that we find

$$a_n^2 = q^{n-1}(1-q^n),$$

which are indeed the recurrence coefficients as given in [12, §3.28]. So for these orthogonal polynomials the recurrence coefficients can be found immediately from the structure relation (5.4). Observe that the a_n^2 tend to zero exponentially fast and that

$$\lim_{n\to\infty} a_n^2 / q^{n-1} = 1,$$

and

$$\lim_{q\to 1} \frac{a_n^2}{1-q^2} = \frac{n}{2},$$

and the latter are the recurrence coefficients (2.8) for the Hermite polynomials ($\rho = 0$).

5.2 Discrete q-Freud polynomials

A q-analog of the Freud polynomials with weight $\exp(-x^4)$ can be obtained by taking the weight $w(x) = (q^4x^4;q^4)_\infty = E_{q^4}(-q^4x^4)$ on the exponential lattice. Observe that $w(\sqrt[4]{1-q^4}x) \to \exp(-x^4)$ as $q \to 1$. This weight satisfies

$$(1-x^4)w(x) = w(x/q), \tag{5.5}$$

and the structure relation for these semi-classical polynomials is:

Lemma 5.2. *The orthonormal polynomials for which*

$$\int_{-1}^{1} p_n(x)p_m(x)(x^4q^4;q^4)_\infty \, d_qx = \delta_{m,n},$$

satisfy

$$D_qp_n(x) = \frac{B_n}{1-q}\, p_{n-1}(x) + \frac{A_n}{1-q}\, p_{n-3}(x), \tag{5.6}$$

with

$$A_n = \frac{a_na_{n-1}a_{n-2}}{q^{n-3}} \tag{5.7}$$

$$B_n = \frac{a_n}{q^{n-1}} \left(\sum_{j=1}^{n+1} a_j^2 - q^2 \sum_{j=1}^{n-2} a_j^2 \right). \tag{5.8}$$

Proof. Expanding D_qp_n into a Fourier series gives

$$D_qp_n(x) = \sum_{j=0}^{n-1} a_{j,n}p_j(x),$$

with

$$a_{j,n} = \int_{-1}^{1} D_qp_n(x)\; p_j(x)w(x)\, d_qx.$$

Again $a_{j,n} = 0$ whenever $n-j$ is even. When $n-j$ is odd then, as in the proof of Lemma 5.1, we have

$$a_{j,n} = \int_{-1}^{1} p_n(x)\frac{p_j(x/q)-p_j(x)}{x(q-1)} w(x)\, d_qx + \frac{1}{1-q}\int_{-1}^{1} x^3p_n(x)p_j(x/q)w(x)\, d_qx,$$

where we have now used the Pearson equation (5.5). Again the first integral on the right vanishes because of orthogonality. The second integral only gives a contribution when $j=n-1$ or $j=n-3$. For $j=n-3$ we have

$$a_{n-3,n} = \frac{1}{1-q}\int_{-1}^{1} x^3p_n(x)p_{n-3}(x)w(x)\, d_qx,$$

and since

$$x^3p_{n-3}(x/q) = \frac{\gamma_{n-3}}{\gamma_n}q^{-n+3}p_n(x) + \text{ lower degree terms}$$

we easily find

$$a_{n-3,n} = \frac{a_n a_{n-1} a_{n-2}}{(1-q)q^{n-3}},$$

which gives (5.7). For $j = n-1$ we have

$$a_{n-1,n} = \frac{1}{1-q} \int_{-1}^{1} x^3 p_n(x) p_{n-1}(x) w(x) \, d_q x,$$

and if we write

$$x^3 p_{n-1}(x/q) = A_{n+2} p_{n+2}(x) + B_n p_n(x) + \text{ lower degree terms,} \tag{5.9}$$

then the orthonormality gives

$$a_{n-1,n} = \frac{B_n}{1-q}.$$

If we compare coefficients of x^n in (5.9) then

$$\delta_{n-1} q^{-n+3} = A_{n+2} \delta_{n+2} + B_n \gamma_n,$$

where $p_n(x) = \gamma_n x^n + \delta_n x^{n-2} + \cdots$, so that using (5.7) gives

$$B_n = \frac{a_n}{q^{n-1}} \left(q^2 \frac{\delta_{n-1}}{\gamma_{n-1}} - \frac{\delta_{n+2}}{\gamma_{n+2}} \right).$$

If we compare coefficients of x^{n-1} in the recurrence relation (5.2) then

$$\delta_n = a_{n+1} \delta_{n+1} + a_n \gamma_{n-1},$$

from which one easily finds

$$\frac{\delta_{n+1}}{\gamma_{n+1}} - \frac{\delta_n}{\gamma_n} = -a_n^2,$$

which gives

$$\frac{\delta_n}{\gamma_n} = -\sum_{j=1}^{n-1} a_j^2, \tag{5.10}$$

and using this in the formula for B_n gives the desired expression (5.8). □

If we compare coefficients of x^{n-1} in the structure relation (5.6) then

$$\gamma_n (1 - q^n) = \gamma_{n-1} B_n. \tag{5.11}$$

Comparing coefficients of x^{n-3} in (5.6) gives

$$\delta_n (1 - q^{n-2}) = \delta_{n-1} B_n + \gamma_{n-3} A_n,$$

which together with (5.11) gives

$$A_n = \frac{\gamma_n}{\gamma_{n-3}} \left(\frac{\delta_n}{\gamma_n} (1 - q^{n-2}) - \frac{\delta_{n-1}}{\gamma_{n-1}} (1 - q^n) \right).$$

Together with (5.7) and (5.10) this gives

$$a_n^2 a_{n-1}^2 a_{n-2}^2 = q^{n-3}\left(q^{n-2}(1-q^2)\sum_{j=1}^{n-2} a_j^2 - (1-q^{n-2})a_{n-1}^2\right). \tag{5.12}$$

On the other hand, if we compare (5.11) with (5.8) then we find

$$\frac{1-q^n}{a_n} = \frac{a_n}{q^{n-1}}\left(\sum_{j=1}^{n+1} a_j^2 - q^2\sum_{j=1}^{n-2} a_j^2\right),$$

which can be written as

$$q^{n-1}(1-q^n) = a_n^2\left(a_{n+1}^2 + a_n^2 + a_{n-1}^2 + (1-q^2)\sum_{j=1}^{n-2} a_j^2\right). \tag{5.13}$$

If we take (5.12) with the index n raised by one, then we can find

$$(1-q^2)\sum_{j=1}^{n-2} a_j^2 = q^{-2n+3}a_{n+1}^2 a_n^2 a_{n-1}^2 - (1-q^2)a_{n-1}^2 - (1-q^{-n+1})a_n^2,$$

and if we insert this in (5.13) then we find the second order non-linear equation

$$q^{n-1}(1-q^n) = a_n^2\left(a_{n+1}^2 + q^{-n+1}a_n^2 + q^2 a_{n-1}^2 + q^{-2n+3}a_{n+1}^2 a_n^2 a_{n-1}^2\right). \tag{5.14}$$

We claim that this equation is a q-deformation of the discrete Painlevé I equation. Indeed, if we take $x_n = a_n^2/\sqrt{1-q^4}$ then

$$q^{n-1}\frac{1-q^n}{1-q^4} = x_n\left(x_{n+1} + q^{-n+1}x_n + q^2 x_{n-1} + (1-q^4)q^{-2n+3}x_{n+1}x_n x_{n-1}\right),$$

which for $q \to 1$ converges to

$$\frac{n}{4} = x_n(x_{n+1} + x_n + x_{n-1}),$$

which is the discrete Painlevé I equation (2.12) for Freud polynomials (with $\rho = 0$). If we put $a_n^2 = q^{n-1}y_n$ then (5.14) can be rewritten as

$$q^n(y_{n+1}y_n + 1)(y_{n-1}y_n + 1) = 1 - y_n^2. \tag{5.15}$$

This could therefore be called a q-discrete Painlevé I equation (q-$\mathrm{P_I}$).

We can easily find the asymptotic behavior as $n \to \infty$. First observe that from (5.14) we find the upper bound

$$q^{-n+1}a_n^4 \leq q^{n-1}(1-q^n),$$

so that $a_n^4 \leq q^{2n-2}(1-q^n)$, and a_n tends to zero as $n \to \infty$. Let $A = \limsup_{n\to\infty} a_n^2/q^{n-1}$, then if we take a such that a_n^2/q^{n-1} converges to A, equation (5.14) gives $A^2 = 1$. A similar reasoning also shows that $B = \liminf_{n\to\infty} a_n^2/q^{n-1}$ is such that $B^2 = 1$. Hence we may conclude that

$$\lim_{n\to\infty} a_n^2/q^{n-1} = 1.$$

The equation (5.15) has the singularity confinement property. Indeed, a singularity occurs for y_{n+1} whenever $y_n = 0$. So if we put $y_n = \epsilon$, then some straightforward calculus gives

$$\begin{aligned} y_n &= \epsilon, \\ y_{n+1} &= q^{-n}(1-q^n)\,\frac{1}{\epsilon} - q^{-n}y_{n-1} + \mathcal{O}(\epsilon), \\ y_{n+2} &= -q^{-n-1}(1-q^n)\,\frac{1}{\epsilon} + y_{n-1}/q + \mathcal{O}(\epsilon), \\ y_{n+3} &= -q^{-2}\frac{1-q^{n+3}}{1-q^n}\,\epsilon + \mathcal{O}(\epsilon^2), \\ y_{n+4} &= q^2\frac{1-q^n}{1-q^{n+3}}\,y_{n-1} + \mathcal{O}(\epsilon). \end{aligned}$$

Hence the singularity is confined to $y_{n+1}, y_{n+2}, y_{n+3}$.

Again the recurrence relation (5.14) or (5.15) is very unstable for computing the recurrence coefficients recursively. One can show [24] that there is again a unique solution of (5.15) with $y_0 = 0$ which is positive for all $n > 0$, and this is the solution for which $y_n = a_n^2/q^{n-1}$. This solution is such that $y_n \to 1$ and

$$y_1 = a_1^2 = \frac{\int_{-1}^{1} x^2(x^4q^4;q^4)_\infty \, d_qx}{\int_{-1}^{1} (x^4q^4;q^4)_\infty \, d_qx} = \frac{(q;q^4)_\infty}{(q^3;q^4)_\infty},$$

where the integrals can be computed using the q-binomial theorem. In Figure 3 we have computed $\log|y_n|$ recursively for $q = 0.9$ with 50 significant digits.



Figure 3: The result of computing $\log|y_n|$ from (5.15) ($q = 0.9$) using 50 significant digits

5.3 Another discrete q-Freud case

If we take the weight $w(x) = (x^2q^2;q^2)_\infty(cx^2q^2;q^2)_\infty$, with $c \leq 1$, then w is positive on the q-lattice and it satisfies the Pearson equation

$$(1-x^2)(1-cx^2)w(x) = w(x/q). \tag{5.16}$$

If $c = -1 + a\sqrt{1-q^4}$ then $w(\sqrt[4]{1-q^4}\,x) \to \exp(-x^4 - 2ax^2)$ so that this gives us a q-deformation of the Freud weight $\exp(-x^4 - 2ax^2)$. Observe that $c = -1$ gives the discrete q-Freud polynomials considered in the previous section and $c = 0$ gives the discrete q-Hermite I polynomials.

Lemma 5.3. *The structure relation for the orthonormal q-polynomials with weight $w(x) = (x^2q^2;q^2)_\infty(cx^2q^2;q^2)_\infty$ on the q-lattice $\{\pm q^n, n \in \mathbb{N}\}$ is*

$$D_q p_n(x) = \frac{\widehat{B}_n}{1-q}p_{n-1}(x) + \frac{\widehat{A}_n}{1-q}p_{n-3}(x), \tag{5.17}$$

where

$$\begin{aligned} \widehat{A}_n &= -c\frac{a_na_{n-1}a_{n-2}}{q^{n-3}} && (5.18)\\ \widehat{B}_n &= -c\frac{a_n}{q^{n-1}}\left(\sum_{j=1}^{n+1} a_j^2 - q^2\sum_{j=1}^{n-2} a_j^2 - \frac{1+c}{c}\right). && (5.19) \end{aligned}$$

Proof. The proof is a straightforward copy of the proof of Lemma 5.2, except that one uses the Pearson equation (5.16). The Pearson equation contains the quartic polynomial $(1-x^2)(1-cx^2) = 1-(1+c)x^2 + x^4$ so that one ends up with integrals of the form

$$\begin{aligned} \widehat{A}_n &= \int_{-1}^{1} [(1+c)x - cx^3]p_n(x)p_{n-3}(x)w(x)\,d_qx = -cA_n \\ \widehat{B}_n &= \int_{-1}^{1} [(1+c)x - cx^3]p_n(x)p_{n-1}(x)w(x)\,d_qx = -cB_n + (1+c)\frac{a_n}{q^{n-1}}, \end{aligned}$$

where A_n and B_n are given by (5.7)–(5.8). □

Reasoning in the same way as in the previous section, i.e., comparing the coefficients of x^{n-1} and x^{n-3} in (5.6), one arrives at

$$q^{n-1}(1-q^n) = -ca_n^2\left(a_{n+1}^2 + q^{-n+1}a_n^2 + q^2a_{n-1}^2 - \frac{1+c}{c} - cq^{-2n+3}a_{n+1}^2a_n^2a_{n-1}^2\right). \tag{5.20}$$

If we put $a_n^2 = q^{n-1}y_n$, then this can be rewritten as

$$(1-y_n)(1-cy_n) = q^n(cy_{n+1}y_n - 1)(cy_{n-1}y_n - 1), \tag{5.21}$$

which is a more general form of the q-discrete Painlevé I equation in (5.15).

Appendix

Several discrete Painlevé equations have appeared in the literature, and the list is certainly longer than the six Painlevé differential equations. Sakai [21] made a classification in terms of rational surfaces associated with affine root systems and the most general (elliptic) discrete Painlevé equation is related with the affine Weyl group symmetry of type E_8. Sakai's classification does not give explicit expressions for the discrete Painlevé equations. A few important discrete Painlevé equations are listed below. The list was compiled by Peter Clarkson and I thank him for his permission to present it in this paper.

A.1 Discrete Painlevé equations

d-P_{I}
$$x_{n+1} + x_n + x_{n-1} = \frac{z_n + \gamma(-1)^n}{x_n} + \delta$$

d-P_{II}
$$x_{n+1} + x_{n-1} = \frac{x_n z_n + \gamma}{1 - x_n^2}$$

d-P_{IV}
$$(x_{n+1} + x_n)(x_n + x_{n-1}) = \frac{(x_n^2 - \kappa^2)(x_n^2 - \mu^2)}{(x_n + z_n)^2 - \gamma^2}$$

d-P_{V}
$$\frac{(x_{n+1} + x_n - z_{n+1} - z_n)(x_n + x_{n-1} - z_n - z_{n-1})}{(x_{n+1} + x_n)(x_n + x_{n-1})} = \frac{[(x_n - z_n)^2 - \alpha^2][(x_n - z_n)^2 - \beta^2]}{(x_n - \gamma^2)(x_n - \delta^2)}$$

where $z_n = \alpha n + \beta$ and α, β, γ, δ, κ, μ are constants.

A.2 q-discrete Painlevé equations

q-P_{II}
$$(x_{n+1}x_n - 1)(x_n x_{n-1} - 1) = \frac{\lambda_n \lambda_{n-1} x_n}{\alpha^2(x_n - \alpha\lambda_n)}$$

q-P_{II}
$$x_{n+1}x_{n-1} = \alpha\lambda_n \frac{\lambda_n + x_n}{x_n(x_n - 1)}$$

q-$\mathrm{P}_{\mathrm{III}}$
$$x_{n+1}x_{n-1} = \frac{(x_n + \alpha)(x_n + \beta)}{(\gamma\lambda_n x_n + 1)(\delta\lambda_n x_n + 1)}$$

q-P_{IV}
$$(x_{n+1}x_n - 1)(x_n x_{n-1} - 1) = \frac{\gamma\delta(x_n + \alpha)(x_n + 1/\alpha)(x_n + \beta)(x_n + 1/\beta)}{(\gamma\lambda_n x_n + 1)(\delta\lambda_n x_n + 1)}$$

q-P_{V}
$$(x_{n+1}x_n - 1)(x_n x_{n-1} - 1) = \frac{\gamma\delta\lambda_n^2(x_n - \alpha)(x_n - 1/\alpha)(x_n - \beta)(x_n - 1/\beta)}{(x_n - \gamma\lambda_n)(x_n - \delta\lambda_n)}$$

q-P_{VI}
$$\begin{aligned} &\frac{(x_n x_{n+1} - \lambda_n\lambda_{n+1})(x_n x_{n-1} - \lambda_n\lambda_{n-1})}{(x_n x_{n+1} - 1)(x_n x_{n-1} - 1)} \\ &\qquad = \frac{(x_n - \alpha\lambda_n)(x_n - \lambda_n/\alpha)(x_n - \beta\lambda_n)(x_n - \lambda_n/\beta)}{(x_n - \gamma)(x_n - 1/\gamma)(x_n - \delta)(x_n - 1/\delta)} \\ &\text{with} \qquad a + b + c + d = 0, \quad p + q + r + s = 0 \end{aligned}$$

where $\lambda_n = \lambda_0 q^n$ and α, β, γ and δ are constants.

A.3 Asymmetric discrete Painlevé equations

$$x_{n+1}=\frac{f_1(y_n)+x_nf_2(y_n)}{f_3(y_n)+x_nf_4(y_n)} \qquad y_{n-1}=\frac{g_1(x_n)+y_ng_2(x_n)}{g_3(x_n)+y_ng_4(y_n)}$$

α-d-P$_{\mathrm{I}}$

$$x_{n+1}+x_n+y_n=\delta+\frac{z_n-\gamma}{y_n}$$

$$y_n+y_{n-1}+x_n=\delta+\frac{z_{n+1/2}+\gamma}{x_n}$$

α-d-P$_{\mathrm{II}}$

$$x_{n+1}+x_n=\frac{2(y_nz_n+\gamma)}{1-y_n^2}$$

$$y_n+y_{n-1}=\frac{2(x_nz_{n+1/2}-\delta)}{1-x_n^2}$$

α-d-P$_{\mathrm{III}}$

$$x_{n+1}x_n=\frac{(y_n-q^na)(y_n-q^nb)}{(y_n-c)(y_n-d)} \qquad y_ny_{n-1}=\frac{(x_n-q^n\alpha)(x_n-q^n\beta)}{(x_n-\gamma)(x_n-\delta)} \qquad \left(\frac{\alpha\beta}{\gamma\delta}=q\frac{ab}{cd}\right)$$

α-d-P$_{\mathrm{IV}}$

$$(y_n+x_n)(x_{n+1}+y_n)=\frac{(y_n-a)(y_n-b)(y_n-c)(y_n-d)}{(y_n+\gamma-z_n)(y_n-\gamma-z_n)}$$

$$(y_n+x_n)(x_n+y_{n-1})=\frac{(x_n+a)(x_n+b)(x_n+c)(x_n+d)}{(x_n+\delta-z_{n+1/2})(x_n-\delta-z_{n+1/2})}$$

with $\qquad a+b+c+d=0$

α-d-P$_{\mathrm{V}}$

$$\frac{(y_n+x_{n+1}-z_n-z_{n+1})(x_{n+1}+y_n-z_n-z_{n-1})}{(y_n+x_n)(x_{n+1}+y_n)}$$

$$=\frac{(y_n-z_n-a)(y_n-z_n-b)(y_n-z_n-c)(y_n-z_n-d)}{(y_n-p)(y_n-q)(y_n-r)(y_n-s)}$$

$$\frac{(y_n+x_{n+1}-z_n-z_{n-1/2})(x_{n+1}+y_n-z_{n-1/2}-z_{n-1})}{(y_n+x_n)(x_n+y_{n-1})}$$

$$=\frac{(x_n-z_{n-1/2}+a)(x_n-z_{n-1/2}+b)(x_n-z_{n-1/2}+c)(x_n-z_{n-1/2}+d)}{(x_n+p)(x_n+q)(x_n+r)(x_n+s)}$$

α-q-P$_{\mathrm{V}}$

$$(x_ny_n-1)(x_{n-1}y_n-1)=q^{2n}\,\frac{(y_n-a)(y_n-b)(y_n-c)(y_n-d)}{(q^n-\kappa y_n)(q^n-y_n/\kappa)}$$

$$(x_ny_n-1)(x_ny_{n+1}-1)=q^{2n+1}\,\frac{(x_n-1/a)(x_n-1/b)(x_n-1/c)(x_n-1/d)}{(q^{n+1/2}-\mu y_n)(q^{n+1/2}-y_n/\mu)}$$

with $\qquad abcd=0$

α-d-P$_{\mathrm{VI}}$

$$\begin{cases} x_nx_{n+1}=\dfrac{\beta_3\beta_4(y_n-q^n\alpha_1)(y_n-q^n\alpha_2)}{(y_n-\alpha_3)(y_n-\alpha_4)} \\ y_ny_{n-1}=\dfrac{\alpha_3\alpha_4(x_n-q^n\beta_1)(x_n-q^n\beta_2)}{(x_n-\beta_3)(x_n-\beta_4)} \end{cases} \qquad \text{with} \qquad \frac{\alpha_1\alpha_2}{\alpha_3\alpha_4}=q\frac{\beta_1\beta_2}{\beta_3\beta_4}$$

where $z_n = \alpha n + \beta$ and a, b, c, d, p, q, r, s, α, β, γ and δ are constants.

A.4 Alternative discrete Painlevé equations

a-d-$\mathrm{P_I}$

$$x_{n+1} + x_n + x_{n-1} = \frac{z_n + \gamma(-1)^n}{x_n} + \mu$$

$$\frac{z_n}{x_{n+1} + x_n} + \frac{z_{n-1}}{x_n + x_{n-1}} = -x_n^2 + \gamma$$

$$x_{n+1} + x_{n-1} = \frac{z_n}{x_n} + \frac{\gamma}{x_n^2}$$

$$x_{n+1} + x_{n-1} = \frac{z_n}{x_n} + \gamma$$

$$x_{n+1}x_{n-1} = \frac{\exp(z_n)}{x_n} + \frac{\gamma}{x_n^2}$$

a-d-$\mathrm{P_{II}}$

$$x_{n+1} + x_{n-1} = \frac{x_n z_n + \gamma}{1 - x_n^2}$$

$$\frac{z_n}{x_{n+1}x_n + 1} + \frac{z_{n-1}}{x_n x_{n-1} + 1} = -x_n + \frac{1}{x_n} + z_n + \gamma$$

a-d-$\mathrm{P_V}$

$$\frac{(x_{n+1} + x_n - 2z_{n+1})(x_n + x_{n-1} - 2z_n)}{(x_{n+1} + x_n)(x_n + x_{n-1})} = \frac{(x_n - z_{n+1/2})^2 - \kappa^2}{(x_n - \gamma)^2}$$

$$\frac{z_{n+1/2}}{1 - x_n x_{n+1}} + \frac{z_{n-1/2}}{1 - x_n x_{n-1}} = \mu + z_n + \frac{\kappa x_n}{(1 + x_n)^2} + \frac{1 - x_n}{1 + x_n}\left[\tfrac{1}{2}z_n + (-1)^n\gamma\right]$$

where $z_n = \alpha n + \beta$ and α, β, γ and δ are constants.

A.5 Other discrete Painlevé equations

$$\begin{array}{lll}
\text{d-P}_{\rm I} & x_{n+1}+x_n=\dfrac{y_nz_n+\gamma}{y_n^2}, & y_n+y_{n-1}=\dfrac{x_nz_{n+1/2}+\delta}{x_n^2} \\[2ex]
\text{d-P}_{\rm II} & x_{n+1}+x_n=\dfrac{y_nz_n+\gamma}{y_n^2-\mu^2}, & y_n+y_{n-1}=\dfrac{x_nz_{n+1/2}+\delta}{x_n^2-\mu^2} \\[2ex]
\text{d-P}_{\rm IV} & x_nx_{n-1}=\dfrac{(y_n+z_n-a)(y_n+z_n-b)}{y_n^2-\gamma^2}, & y_n+y_{n+1}=-\dfrac{z_{n+1/2}+c}{x_n\gamma+1}-\dfrac{z_{n+1/2}+d}{x_n/\gamma+1} \\
 & \quad\text{with}\quad a+b+c+d=0 & \\[3ex]
\text{d-P}_{\rm IV} & x_nx_{n-1}=\dfrac{a(y_n+z_n-b)}{y_n^2-\gamma^2}, & y_n+y_{n+1}=\dfrac{c}{x_n}+\dfrac{z_{n+1/2}+d}{x_n-1} \\[2ex]
\text{d-P}_{\rm V} & x_n+x_{n-1}=\dfrac{z_n+\mu}{1+y_n/t}+\dfrac{z_n-\mu}{1+ty_n}, & y_ny_{n+1}=\dfrac{(x_n-z_{n+1/2})^2-\kappa^2}{x_n^2-\gamma^2} \\[2ex]
 & x_n+x_{n-1}=\dfrac{\gamma}{1+y_n}+\dfrac{z_n+\delta}{1-y_n}, & y_ny_{n+1}=\gamma\,\dfrac{x_n-z_n}{x_n^2-\mu^2}
\end{array}$$

where $z_n=\alpha n+\beta$ and a, b, c, d, p, q, r, s, α, β, γ, δ, κ and μ are constants.

References

[1] J. Baik, *Riemann-Hilbert problems for last passage percolation*, in "Recent Developments in Integrable Systems and Riemann-Hilbert Problems" (K. McLaughlin, X. Zhou, eds.), Contemporary Mathematics **326** (2003), pp. 1–21.

[2] J. Baik, P. Deift, K. Johansson, *On the distribution of the length of the longest increasing subsequence of random permutations*, J. Amer. Math. Soc. **12** (1999), 1119–1178.

[3] T. S. Chihara, *On quasi-orthogonal polynomials*, Proc. Amer. Math. Soc. **8** (1957), 765–767.

[4] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Mathematics and its Applications **13**, Gordon and Breach, New York, 1978.

[5] R. Conte (editor), *The Painlevé Property. One Century Later*, CRM Series in Mathematical Physics, Springer-Verlag, New York, 1999.

[6] C. Cresswell, N. Joshi, *The discrete Painlevé I hierarchy*, in 'Symmetries and Integrability of Difference Equations' (Canterbury, UK, July 1-5, 1996), Lond. Math. Soc. Lect. Notes Ser. **255**, (P.A. Clarkson et al., eds.), Cambridge University Press (1999), pp. 197–205.

[7] G. Freud, *On the coefficients in the recursion formulae of orthogonal polynomials*, Proc. Roy. Irish Acad. Sect. A **76** (1976), no. 1, 1–6.

[8] B. Grammaticos, F. W. Nijhoff, A. Ramani, *Discrete Painlevé equations*, in [5], pp. 413–516.

[9] M. Hisakado, *Unitary matrix models and Painlevé III*, Mod. Phys. Letters **A11** (1996), 3001–3010.

[10] M. N. Hounkonnou, C. Hounga, A. Ronveaux, *Discrete semi-classical orthogonal polynomials: generalized Charlier*, J. Comput. Appl. Math. **114** (2000), 361–366.

[11] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, Encyclopedia of Mathematics and its Applications **98**, Cambridge University Press, 2005.

[12] R. Koekoek, R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue*, Reports of the faculty of Technical Mathematics and Informatics no. 98-17, Delft University of Technology, 1998.

[13] J. S. Lew, D. A. Quarles, *Nonnegative solutions of a nonlinear recurrence*, J. Approx. Theory **38** (1983), no. 4, 357–379.

[14] D. S. Lubinsky, H. N. Mhaskar, E. B. Saff, *A proof of Freud's conjecture for exponential weights*, Constr. Approx. **4** (1988), no. 1, 65–83.

[15] A. P. Magnus, *A proof of Freud's conjecture about the orthogonal polynomials related to* $|x|^\rho \exp(-x^{2m})$*, for integer* m, in 'Orthogonal Polynomials and Applications' (Bar-le-Duc, 1984), Lecture Notes in Mathematics **1171**, Springer, Berlin, 1985, pp. 362–372.

[16] A. P. Magnus, *On Freud's equations for exponential weights*, J. Approx. Theory **46** (1986), no. 1, 65–99.

[17] A. P. Magnus, *Freud's equations for orthogonal polynomials as discrete Painlevé equations*, in 'Symmetries and Integrability of Difference Equations' (Canterbury, 1996), London Math. Soc. Lecture Note Series **255**, Cambridge University Press, Cambridge, 1999, pp. 228–243.

[18] A. Máté, P. Nevai, T. Zaslavsky, *Asymptotic expansions of ratios of coefficients of orthogonal polynomials with exponential weights* Trans. Amer. Math. Soc. **287** (1985), no. 2, 495–505.

[19] F. W. Nijhoff, *On a q-deformation of the discrete Painlevé I equation and q-orthogonal polynomials*, Lett. Math. Phys. **30** (1994), 327–336.

[20] V. Periwal, D. Shevitz, *Unitary-matrix models as exactly solvable string theories*, Phys. Rev. Letters **64** (1990), 1326–1329.

[21] H. Sakai, *Rational surfaces associated with affine root systems and geometry of the Painlevé equations*, Commun. Math. Phys. **220** (2001), no. 1, 165–229.

[22] B. Simon, Orthogonal Polynomials on the Unit Circle, Amer. Math. Soc. Colloq. Publ. **54**, Part 1 and Part 2, Amer. Math. Soc., 2005.

[23] C. A. Tracy, H. Widom, *Random unitary matrices, permutations and Painlevé*, Commun. Math. Phys. **207** (1999), 665–685.

[24] W. Van Assche, *Unicity of certain solutions of the discrete Painlevé II equation*, manuscript

[25] W. Van Assche, M. Foupouagnigni, *Analysis of non-linear recurrence relations for the recurrence coefficients of generalized Charlier polynomials*, J. Nonlinear Math. Phys. **10** (2003), suppl. 2, 231–237.

Katholieke Universiteit Leuven
Department of Mathematics
Celestijnenlaan 200 B
B-3001 Leuven
BELGIUM
`walter@wis.kuleuven.be`